



# **Agromet Advisory Bulletin**

Date : 27-08-2024

Weather Forecast of District MEDINIPUR-EAST(West Bengal) Issued On : 2024-08-27(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-08-28	2024-08-29	2024-08-30	2024-08-31	2024-09-01
Rainfall(mm)	20.0	15.0	25.0	17.0	12.0
Tmax(°C)	32.0	33.0	33.0	33.0	34.0
Tmin(°C)	27.0	27.0	27.0	27.0	28.0
RH-I(%)	86	85	88	89	90
RH-II(%)	64	66	67	72	64
Wind Speed(kmph)	12	10	17	14	11
Wind Direction(Degree)	169	154	90	135	120
Cloud Cover(Octa)	8	7	4	5	6

#### Weather Summary/Alert:

Heavy rainfall is predicted on 28th and 30th August and moderate rainfall is predicted in the remaining days. The sky will be partly to mainly cloudy in next 5 days. Maximum temperature is predicted around 32 to 34 degrees and minimum temperature is predicted around 27 to 28 degrees in next 5 days. Maximum humidity is forecasted around 85-90% and minimum humidity 64-72%. Wind speed will be 10-14 km/hr and the predominant wind direction will be South Easterly to South Westerly.

#### **General Advisory:**

As per extended range forecast deficit in rainfall was recorded from 8.8.24 to 21.8.24. Based on MME forecast above normal rainfall is predicted from 23.8.24 to 29.8.24 and normal rainfall is predicted from 30.8.24 to 5.9.24. Based on MME forecast Tmax will be normal in Week 1 and week 2. For Tmin, it will show normal in Week 1 and week 2. ------ Advisory: Forecast for moderate rainfall, prediction is there for rising maximum temp and humidity , chance of huge sheath blight and sheath rot attack in aman paddy field. If attack found apply Validamycine (COntaf plus) @ 1ml per liter of water. Spraying must be done in bund area also.

#### **SMS Advisory:**

Forecast for moderate rainfall, prediction is there for rising maximum temp and humidity, chance of huge sheath blight and sheath rot attack in aman paddy field. If attack found apply Validamycine

### **Crop Specific Advisory:**

Crop (Stage)	Crop Specific Advisory		
RICE (Tillering)	South 24 Parganas Aman paddy Early transplanted paddy in tillering stage		
	and Late transplanting paddy in vegetative stage. As more than 100mm		

Crop (Stage)	Crop Specific Advisory		
	rainfall is received deep water stagnation restrict application of first top		
	dressing and herbicide application. Seedlings damaged due to deep water		
	stagnation • No spraying for weed control should be applied on watered field.		
	• Where newly transplanted main field was found completely submerged		
	replace the damaged seedlings with healthy one or tiller from early		
	transplanted paddy field. • Saved seedlings can be used for re-transplanting		
	till no node will form • Re transplanting can be continued up to 1st Week of		
	September Purba Medinipur Aman paddy Tillering stage Observation should		
	be done during morning hours to inspect pest attack condition • Where basal		
	dose cannot be given due to water stagnation apply DAP (granular) and		
	Potash fertilizer@4-5kg per bigha. • If leaf roller or stem borer moth found		
	during inspection only then Chlorothalonyprol can be applied.		

## Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory		
BRINJAL (Grand growth)	Brinjal, Tomato, Chilli Tender Seedling • Drain out of excess water • No spraying or pesticide should be applied in next 2-3 days. • After removal of water apply Diethane M-45@2.5g/liter of water for preventing wilting in sunny days.		
BETELVINE (Grand growth)	Betel vine GrowingWater stagnation caused rotting of vine and leaves • Immediately drain out water and pluck the affected leaf from field • Apply 0.5% Bordeaux mixture • Arrange proper drainage facility as moderate rainfall is predicted.		