## BIDHAN CHANDRA KRISHI VISWAVIDYALAYA MEDIUM-RANGE WEATHER FORECAST AND AGRO-ADVISORY (Agromet Advisory Services, IMD, MoES)

Directorate of Research, B.C.K.V., Kalyani.



E-mail: <u>aaskalyani@gmail.com</u>, <u>bhowmickmahasweta2012@gmail.com</u>

From <u>Kalyani (</u>Nadia, WB)

Weather Forecast for Burdwan District

## FORECAST PERIOD: 25th April to 29th April, 2018

## Weather forecast for next five days

- Clear sky is expected during first 3 days and partly cloudly sky is expected during last 2 days.
- Light rain during next few days is expected except 25th April(no rain).Total rainfall will be 31mm.
- ➤ Wind flow mainly from S & SE direction with 10-21 km/hr. speed during next 5 days is expected.
- Maximum relative humidity is expected to be around 72-%-86% and minimum relative humidity is expected to be around 9% - 39%.
- ➤ Maximum temperature is expected to be around 33°C -36°C and minimum temperature is expected to be around 24 °C -26 °C

| Crop                            | Stage                     | Advisory  |
|---------------------------------|---------------------------|---|
| Cucumber                        | Fruiting                  | Mulching may be practiced. Collection and destruction of fruit fly affected fruits need to be done.   |
| Green<br>manuring<br>crop       | Sowing                    | Sowing of green manuring crop like cluster bean, dhaincha etc may be<br>advised where harvesting of rabi crop is over.<br>Ensure proper moisture for proper germination and growth of crop.   |
| Mango                           | Fruit<br>setting          | In mango orchard, to prevent shedding of new fruits due to high temperature apply irrigation and spray NAA@ 5ml/10 lt. of water.  |
| Onion,<br>Garlic<br>&<br>Tomato | Harvesting                | Keeping in view of forecasted high temperature, farmers are advised<br>for harvesting of mature vegetables at morning and evening.  |
| Jute                            | Seed-sowing<br>& Seedling | Treat the seeds with carbendizm 50WP @ 1g/kg of seed before<br>sowing. Apply Trichoderma viride mixing with organic manure @<br>5g/kg of organic manure in field soil. In case of rotting of seedlings<br>i.e.seedling blight apply carbendazim 50 WP @1g/lit. of water or Cu-<br>hydroxide 75 WP@ 2g/lit.of water. |

## **Agrometeorological Advisory**