



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
**Bidhan Chandra Krishi Viswavidyalaya**  
 Mohanpur, Dist- Nadia, West Bengal



## Agromet Advisory Bulletin

Date : 26-11-2021

Weather Forecast of HOOGLHY(West Bengal) Issued On : 2021-11-26(Valid Till 08:30 IST of the next 5 days)

Parameter	2021-11-27	2021-11-28	2021-11-29	2021-11-30	2021-12-01
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	29.0	29.0	28.0	28.0	28.0
Tmin(°C)	19.0	18.0	18.0	18.0	18.0
RH-I(%)	61	60	59	62	65
RH-II(%)	37	37	33	33	35
Wind Speed(kmph)	8.2	9.2	10.5	9.6	11.4
Wind Direction(Degree)	27	22	338	338	207
Cloud Cover(Octa)	0	4	8	5	7

### Weather Summary/Alert:

No rainfall is likely to occur during next few days. Partly cloudy to mainly cloudy sky is expected on next few days . Wind speed is expected to be 8-12 km / hour. Max. RH is expected to be 59% to 65% . Maximum temperature is expected to be around 28.0-29.0 deg C and minimum temperature is expected to be around 18.0-19.0 deg C.

### General Advisory:

Please download Mausam , Meghdoot & Damini App at your smart phone to get weather forecast and agromet advisory .

### SMS Advisory:

No rain is expected , spraying , fertilizer application , harvesting may be done .

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Continue rice harvesting . After drying & threshing ,grains should be properly dried to right moisture level. Don't burn the residue of the crop.

### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
-------------------------	--------------------------------

<b>Horticulture(Varieties)</b>	<b>Horticulture Specific Advisory</b>
POTATO	Complete planting of potato . N:P:K dose 150 : 150 : 150 kg / ha may give the good result . The crop matures in 80-90 days. Medium high land , sandy loam to loamy soil , slightly acidic to neutral ph is ideal for potato cultivation. One deep ploughing , 2-3 cross & shallow ploughing make the soil well pulverized , loose , & friable with good tilth . 10- 15 ton well rotten organic manure must be mixed during first ploughing . Apply phorate 10 G @ 15 Kg/ha to control soil borne insect. Collect certified seed tuber from Govt. approved organization . 25-45 g seed tuber must be treated with trichoderma viridi or pseudomonas fluorescence solution. Recommended dose of N :P:K is 180-200 : 150 : 150 Kg/ha. Nitrogen should be applied in split doses . Zn & B may be applied in case of micronutrient deficiency in the form of ZnSo4 (25 Kg/ha ) and Borax (10 Kg/ha ) at the time of final land preparation