12/17/21, 1:32 PM District Advisory



Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Bidhan Chandra Krishi Viswavidyalaya



Mohanpur, Dist- Nadia, West Bengal

Agromet Advisory Bulletin

Date: 17-12-2021

Weather Forecast of HOOGHLY(West Bengal) Issued On: 2021-12-17(Valid Till 08:30 IST of the next 5 days)

Parameter	2021-12-18	2021-12-19	2021-12-20	2021-12-21	2021-12-22
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	26.0	25.0	23.0	22.0	23.0
Tmin(°C)	15.0	14.0	11.0	11.0	12.0
RH-I(%)	51	58	44	54	56
RH-II(%)	27	31	22	25	32
Wind Speed(kmph)	9.8	14.4	12.8	9.4	9.3
Wind Direction(Degree)	338	338	335	333	333
Cloud Cover(Octa)	2	1	0	0	0

Weather Summary/Alert:

No rainfall is likely to occur during next few days. Clear sky is expected on next few days. Wind speed is expected to be 10-14 km / hour. Max. RH is expected to be 48% to 64%. Maximum temperature is expected to be around 24.4-26.9 deg C and minimum temperature is expected to be around 10.1-14.0 deg C.

General Advisory:

Prepare drainage channels in the fields to immediately drain out the stagnant water from standing crop which is created due to light / moderate/ heavy rainfall of last week's depression rainfall . Apply Blitox @ 4 g/ lt of water in a clear , sunny weather in the morning or afternoon hour. Weeding is recommended with pre emergence herbicide . Stop applying N fertilizer as top dressing at this moment. Download Mausam, Meghdoot & Damini apps at your smart phone .

SMS Advisory:

Direct seeded spinach, radish, carrot etc may be sown as 2.5 months crop in the fields where newly sown Potato or Mustard are totally affected due to water logging condition for recent depression.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
LENTIL	Drain the excess water immediately . For leaf spot , leaf blight, root rot , wilt etc spray Carbendazim (1 g / lt water) and Tebuconazole (1.5 ml per lt of water) may be applied at 1 week interval.

12/17/21, 1:32 PM District Advisory

Crop(Varieties)	Crop Specific Advisory			
INTERNATION	Drain the excess water immediately to avoid water logging. For alternaria blight, stem or root rot spray with Metalaxyl and Mancozeb @ 2.5 g /lt of water.			

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory		
SPINACH	Direct seeded spinach, radish, carrot etc may be sown as 2-2.5 months crop in the fields where newly sown Potato or Mustard are totally affected due to water logging condition for recent depression rainfall.		
POTATO	Drain the excess water immediately.Prophylactic spraying with Mancozeb (for early blight) @ 2.5gm/lt water and Metribuzine @ 10 g/ per drum water (for weed) is recommended in a clear sunny day .		